

Snail fauna of the oldest cemeteries from Riga (Latvia)

JOZEF ŠTEFFEK¹, ARTURS STALAŽS² & EDGARS DREIJERS³

¹Faculty of Ecology and Environmental Science, Technical University of Zvolen, Institute of Forest Ecology, Slovak Academy of Sciences, Štúrova 2, SK-960 53 Zvolen, Slovak Republic; e-mail: steffekjozef@yahoo.com

²Latvia State Institute of Fruit-Growing, Graudu iela 1, Dobeles nov. (Dobeles raj.), LV-3701, Latvia; e-mail: arturs.stalazs@lvai.lv

³University of Latvia, Faculty of Biology, Kronvalda bulvāris 4, Rīga, LV-1010, Latvia; e-mail: helix@inbox.lv

ŠTEFFEK J., STALAŽS A. & DREIJERS E., 2008: Snail fauna of the oldest cemeteries from Riga (Latvia). – Malacologica Bohemoslovaca, 7: 79–80. Online serial at <<http://mollusca.sav.sk>> 29-Sep-2008

We report on the snail fauna from one of the oldest cemeteries from Riga city (Latvia), viz. Lielie kapi and Pokrova kapi. A list of all the snail species recorded from both cemeteries and limited information on the biotope conditions are given. *Ceciloides acicula* (O.F. Müller, 1774) is first recorded from Latvia and was found at a grassland within the tomb area of the Pokrova kapi cemetery.

Key words: Molluscs, land snail, cemeteries, *Ceciloides acicula*, Riga, Latvia

Introduction

Historical cemeteries represent a suitable refuge for snail fauna within an urban area. Their positive influence is mainly due to dense plant and tree (*Tilia* sp., *Acer* sp., *Fraxinus* sp.) cover which form favourable microclimatic and soil conditions. Yet, such biotopes, due to the introduction of exotic plants, also serve as potential vectors for biological invasions of foreign snail and slug species. In this paper, we present a checklist of the snail fauna of two historical cemeteries, viz. Lielie kapi (Grate Cemetery) and Pokrova kapi (Pokrow's Cemetery), in the urban area of Riga, Latvia (50°85' N, 63°13' E).

Study area

The history of the cemetery of Lielie kapi starts from 1773. Originally, until the 18th century, the area was a sandy field. Nowadays, Lielie kapi is a monumental park with many walls, monuments and a few chapels. It was closed as an active cemetery in 1969, and was later transformed to a monumental park. The Pokrova kapi is still in use. Parts of Lielie kapi and Pokrova kapi are separated from the major area of Lielie kapi by the Senču Street. The detached parts of both cemeteries differ from the larger part in a higher cover of shrubs and trees.

The dominant tree species at both cemeteries are lime (*Tilia* sp.), maple (*Acer* sp.) and oak (*Quercus* sp.) together with less abundant *Betula*, *Aesculus* and several *Populus*, *Fraxinus* and *Thuja* trees. Most of the trees have been planted artificially. The dense tree canopy forms dark conditions at both cemeteries, especially at the Pokrova kapi cemetery that has a dense understory of small trees. The herbal layer is dominated by the nitrophilous species *Ae-*

gopodium podagraria, *Urtica dioica* and *Lamium album*. Regular grass cutting is applied in the largest part of the Lielie kapi cemetery.

During the Baltic malacozoological conference, held on the September 21st–24th 2006 in Riga, authors visited the area of cemeteries Lielie kapi and Pokrova kapi. We sampled the (decaying) leaf litter layer at several sites, which were searched for the presence of snails.

Results and discussion

A total of 17 mollusk species was recorded at both cemeteries (Table 1). The highest species richness was recorded at the Pokrova kapi cemetery and in the vicinity of building walls (15 species). The open grassland with intensive grass-cutting were rather poor in recorded species (six species). Two land snail species were abundant in this area, viz. *Arianta arbustorum* and *Trichia hispida*. A high density of the introduced, and invasive, species *Krynckillus melanocephalus*, which is regularly found in the urban area of Riga city, was not found in the study area. This may be caused by the very dry weather conditions during the months before sampling.

Ceciloides acicula (O.F. Müller, 1774) is first recorded from Latvia. The geographical range of *C. acicula* is the W Palaearctic and Central Mediterranean where it can be found in low- as well as upland. This species lives subterraneous. It has been recorded across Europe, common also in the southern part of Scandinavia and in Great Britain (KERNEY et al. 1983, WIKTOR 2004). Until now, the species was not known from Latvia (RUDZITE et al. 1996, 1997, RUDZITE 2000). In Riga, it was found together with the Caucasian slug *Krynckillus melanocephalus*

Table 1. Snail species found in cemeteries Lielie kapi and Pokrova kapi (Riga City, Latvia), C1 - Pokrova kapi and Lielie kapi (smaller part), C2 - Lielie kapi (larger part).

Species	C1	C2	Comments
<i>Arianta arbustorum</i> (Linnaeus, 1758)	+	+	
<i>Arion</i> (<i>Carinarion</i>) sp. juv.	+	+	
<i>Arion silvaticus</i> Lohmander, 1937	+	+	
<i>Ceciloides acicula</i> (O.F. Müller, 1774)	+	+	new for Latvia
<i>Cochlicopa lubrica</i> (O.F. Müller, 1774)	+	+	
<i>Cochlodina laminata</i> (Montagu, 1803)	+	–	
<i>Deroceras reticulatum</i> (O.F. Müller, 1774)	+	+	
<i>Discus rudersatus</i> (W. Hartmann, 1821)	+	–	
<i>Helix pomatia</i> Linnaeus, 1758	–	+	non-native
<i>Krynickyllus melanocephalus</i> Kaleniczenko, 1851	+	+	non-native
<i>Limax maximus</i> Linnaeus, 1758	+	–	non-native
<i>Oxychilus draparnaudi</i> (H. Beck, 1837)	+	+	non-native
<i>Oxychilus translucidus</i> (Mortillet, 1881)	–	+	non-native
<i>Succinea putris</i> (Linnaeus, 1758)	+	–	
<i>Trichia hispida</i> (Linnaeus, 1758)	+	+	
<i>Vallonia costata</i> (O.F. Müller, 1774)	+	+	
<i>Vallonia pulchella</i> (O.F. Müller, 1774)	+	+	
Total 17 sp.			

Explanatory notes: + species present, – species absent

Kaleniczenko, 1851 and *Oxychilus translucidus* (Mortillet, 1853) on a grassland within the tomb area in the Pokrova kapi cemetery (10 shells leg., det. et coll. J. Šteffek, 24 Sep 2006). *K. melanocephalus* was found for the first time by A. Stalažs in 1997 in the National Botanical Gardens of Latvia (Salaspils, Riga district). The species was only recognized in 2000 after dissection by E. Dreijers and A. Wiktor. The first record of *O. translucidus* is from 1999 from greenhouses of the National Botanical Gardens of Latvia (Leg. A. Stalažs) and was identified by anatomical dissection by E. Dreijers after 2000 (E. DREIJERS & A. STALAŽS, unpubl. data). Both species belong to alien molluscs and *K. melanocephalus* appears to be an invasive species, which is known from several localities in Hungary, Bulgaria, Turkey, Poland and Israel (WIKTOR 2004), Latvia (DREIJERS 2003) and Germany (MENG & BÖSSNECK 1999, BÖSSNECK & FELDMANN 2003).

In the Baltic States, *O. translucidus* is one of the three alien species from the genus *Oxychilus* (*O. cellarius*, *O. draparnaudi* and *O. translucidus*) and the species occurs in Latvia (several localities) and Lithuania (one record from Kaunas by A. Stalažs in 2003). Only *O. alliarius* (Miller, 1822) is a native species in the Baltic States (E. DREIJERS & A. STALAŽS, unpubl. data).

Acknowledgements

The study was funded by research grants no. 1/3283/06, 2/6007/26 and 2/5152/25 of the Slovak Grant Agency for Science (VEGA).

References

- BÖSSNECK U. & FELDMANN A., 2003: Zur Ausbreitung von Neozoa im Stadtgebiet von Erfurt am Beispiel der Landschnecken *Cer-nuella neglecta* (Draparnaud, 1805), *Monacha cartusiana* (O.F. Müller, 1774) and *Krynickyllus melanocephalus* Kaleniczenko, 1851 (Mollusca: Gastropoda). – Veröffentlichungen Naturkundu-museum Erfurt, 22: 115–125.
- DREIJERS E., 2003: A history of the investigation and fauna of the slugs (Gastropoda terrestria nuda) of Latvia, p. 33. – In: Second International Conference „Research and conservation of biological diversity in Baltic Region“, book of Abstracts, Daugavpils, 24.–26.04.2003.
- KERENY M.P., CAMERON R.A.D. & JUNGBLUTH J.H., 1983: Die Landschnecken Nord- und Mitteleuropas. – Verlag Paul Parey, Hamburg/Berlin.
- MENG S. & BÖSSNECK U., 1999: *Krynickyllus melanocephalus* Kaleniczenko, 1851 in Deutschland eingeschleppt (Gastropoda: Stylomatophora: Agriolimacidae). – Malk. Abh. Mus. Tirkde. Dresden Bd., 19: 304–309.
- RUDZITE M., 2000: Mācies pazīt – Latvijas zemesgliemeži. – Riga, 147 pp.
- RUDZITE M., PILATE D. & PARELE E., 1996: Latvija sastopamo gli-emju sugu latviskie nosaukumi. – Daba un Muzejs, 6: 85–92.
- RUDZITE M., PILATE D. & PARELE E., 1997: Molluskenfauna Let-lands. Liste der in Lettland vorkommenden Molluskenarten (Gastropoda, Bivalvia). – Mitt. Dtsch. malakozool. Ges., 59: 1–10.
- SCHLESCH H., 1942: Die Land- und Süßwassermollusken Let-lands. – Korrespondenzblatt des Naturforscher-Verein zu Riga, 64, 311 pp.
- WIKTOR A., 2004: Ślimaki lądowe Polski. – Wydawnictwo Man-tis, Olsztyn, 302 pp.